Exclusion criteria was: not willing to participate
Chronically ill with systemic involvement

Predesigned semi structured questionnaire was used to elicit information on general socioeconomic profile, addiction, awareness regarding reproductive sexual health along with awareness on HIV/AIDS and also their health seeking behavior. The opportunity was used to impart them health education and refer appropriately. Anonymity of the respondent was assured.

The data was compiled and analyzed using SPSS software for any statistical significance.

Results: A total of 168 workers were recruited in the survey, with 68% females; 68% were married; 52% within 31–54 years; 56% had substance abuse of which 44% used gutka; 32% felt they were not reproductively healthy out of which only 58% sought medical help in spite of being working in the medical college out of which only 20% went to indigenous college. 24% of those who sought help were non compliant with therapy. Spouses (52%) were key confidantes. Both among females (48%) and males (68%) complained of burning micturition. 18% reported having never heard of HIV/AIDS and benefits of condoms against the disease were unknown to nearly 47%. Unmet need of contraception was nearly 43%. Education of self and spouse was seen to have a strong association with the RSH.

Conclusion: Reproductive and Sexual Health is an identified taboo subject yet neglected in all sections of the society. The study shows dismal levels of awareness and health seeking behavior even in a cohort that works in close proximity with health providers. This hints that other sections might be even worse affected and in need of dire intervention. Counseling and screening of such workforce at regular intervals would be a good solution.

Prevalence of hypertension among private school teachers in Chennai – A cross sectional study

J. Carl Lewis, S. Valarmathi, J. S. Sundar, S. Parameswari S, D. Shantharam

Department of Epidemiology, The Tamilnadu Dr. MGR Medical University, India

Background: Hypertension is reported to be the 4th & 7th contributor to premature death in developed and in developing countries respectively. Recent reports indicate nearly 1 billion adults had hypertension in 2000, it is predicted to increase to 1.56 billion by 2025. Hypertension is considered as an important independent predictor of cardiovascular disease, cerebro-vascular accidents and death.

Objectives: To estimate the prevalence of hypertension and find the associated risk factors among BT (Bachelor of Teaching) and PG (Post Graduate)-assistant teachers.

Design: Cross Sectional Study.

Setting: Private School from Educational Zones of Chennai.

Methods: A sample of 179 teachers participated in the survey. After obtaining informed consent socio-demographic details, food habits, physical-activity, anthropometric measurements and family history were collected. Blood Pressure was measured at 0, 5 & 30 minutes, using digital apparatus and average of three readings was taken as BP of the Individual.

Results: Among 179 teachers males (18%) and females (82%). The prevalence of hypertension: overall-12.8%, BT-assistant-9.3%, PG-assistant-17%. Chi-square-test resulted that there was a significant ($p < 0.05$) association between hypertension and Age, BMI, Physical activity. Backward multiple logistic regression resulted Age, BMI, Food habits, Aerated-drinks are significant risk factors. Hosmer and Lemeshow test for goodness of fit was used to assess the multiple logistic regressions. It resulted in a non-significant value which is an indicative that the model predicts the population fairly well.

Conclusion: Hypertension was high among PG-assistant compared to BT-assistant. Life-style modifications, increased physical activity, diet modification is recommended to reduce emergence, prevalence of hypertension and thereby reducing the risk of cardiovascular disease.

Copyright © 2013, INDIACLEN. Publishing Services by Reed Elsevier India Pvt Ltd. All rights reserved.

<http://dx.doi.org/10.1016/j.cegh.2013.04.003>

Original Article

Profession of barbering: Unexplored issues in HIV/AIDS

Bhola Nath^{a,*}, Ranjeeta Kumari^b, Tanu Midha^c, Narain Das Vaswani^d, Seema Lekhwani^e

* Corresponding author. Tel.: +91 8528463091.

E-mail addresses: bhola_2001@sify.com, bholanath75@yahoo.co.in (B. Nath).

^a Associate Professor, Department of Community Medicine, V.C.S.G. Government Medical College, Srinagar, Uttarakhand 246174, India

^b Assistant Professor, Department of Community Medicine, AIIMS, Rishikesh, India

^c Assistant Professor, Department of Community Medicine, G.S.V.M.M.C., Kanpur, India

^d Associate Professor, Department of Pediatrics, PBDSPGIMS, Rohtak, India

^e Assistant Professor, Department of Biochemistry, PBDSPGIMS, Rohtak, India

ARTICLE INFO

Article history:

Received 18 September 2012

Accepted 27 November 2012

Available online 6 June 2013

Keywords:

Barber

HIV/AIDS

Mode of transmission

ABSTRACT

Background: Barbers are one of the potential high-risk groups due to the increased risk of injuries owing to their profession as well as their exposure to all the sections of the society. Lack of knowledge regarding HIV and appropriate disinfection procedures might provide a fertile ground for propagation of the disease.

Methodology: A cross-sectional survey was conducted in Kanpur, India from August to September, 2009 to assess the awareness and practices of shop barbers (SB) and road side barbers (RSB) regarding HIV.

Results: The mean age group of the barbers was 32.05 ± 11.02 years. Knowledge regarding all the modes of transmission especially blood contact (5.6%) was known to very few RSB. Myths regarding the mode of transmission were rampant. The overall knowledge regarding the disinfection of the instruments (42.8%) and use of new blades (60.6%) was average. Significant differences were observed in the knowledge and practices of the SB and RSB.

Conclusion: Significant gap in knowledge and practices were observed in the barbers specially the RSB. Due to the unique position that the barbers enjoy in the society they can be educated regarding HIV and subsequently used as peer educators for impeding the march of HIV.

Copyright © 2012, INDIACLEN. Publishing Services by Reed Elsevier India Pvt Ltd. All rights reserved.

Key message: The profession of barbering poses a definite risk of transmission of HIV along with other blood borne infections and therefore health education to improve knowledge regarding HIV and methods of prevention to be applied in the barbers' shop need to be disseminated and adhered to.

1. Introduction

National AIDS Control Policy (1999–2004) of India has recognized AIDS as not “merely a public health challenge,” but also a political and social development issue.¹ By doing so, National AIDS Control Organization (NACO) has officially recognized that the epidemic has moved from vulnerable or ‘high risk’ groups to the general population. This has occurred through the various modes of transmission, one of which may be through the cuts and abrasions at the much neglected barbers shop^{2–5} One goal of the policy has therefore been to ensure the safe use of sharp equipments. The barbers shop is a place for the regular visit by people from all the sections of the society including those from high-risk behavior group for HIV like laborers, rickshaw-pullers, truck-driver, migrants and slum dwellers. It therefore serves as a conglomeration place for the transmission of various infections.⁶

Though many studies related to knowledge, attitude and practices and their effect on HIV transmission have being conducted among various sectors of society, little attention has been paid to the profession of hair cutting and the use of sharp equipments in the profession of barbering.

Human immunodeficiency virus (HIV) the agent responsible for causing AIDS is known to be transmitted from one person to another through the use of non-sterile needles, syringes and other skin piercing equipments such as blades and scissors.^{7–9}

Proper sterilization of all such instruments is therefore important in order to prevent HIV transmission. Fortunately, HIV is very sensitive to standard methods of sterilization.^{10,11}

However, barber's shop is a place where there is frequent use of the same blade, trimmers and scissors, often without proper sterilization or disinfection. The use of these sharp instruments may represent an HIV hazard to the general population due to skin piercing injuries.^{12,13}

In India, two types of barber shops are available; one is the road side barber where there is no proper infrastructure and other is an enclosed shop. Hence the present study was carried out to assess the awareness and reported practices of barbers with respect to HIV and also to see if there were any differences based on the type of shop.

2. Methodology

A cross-sectional survey was conducted in Kanpur city, India from August to September, 2009 after obtaining approval from Ethical Committee. A structured questionnaire was used to assess the awareness and practices of barbers regarding HIV. Kanpur is a city of northern India on the Ganges River and is very densely populated. Urban Kanpur has 6 divisions (also known as zones) consisting of 110 wards with a total

population of 2,797,511 according to 2001 census. There are numerous big and small barber shops and a number of road side barbers catering to the population. Two-stage stratified random sampling technique was used. In the first stage, we selected 3 divisions randomly out of 6 divisions in the district. In the second stage, from each division we selected 6 wards randomly out of 18–19 wards in each division according to the list of wards obtained from Nagar Nigam, Kanpur. Finally interviews were conducted with 20 barbers in each ward after obtaining informed consent from them. We selected 15 shop barbers (SB) and 5 road side barbers (RSB) from each ward, due to the differences observed in their proportions. Therefore a total of 360 barbers were included in the study. The selection of the barbers was done non-randomly due to the non-availability of the list of barbers. The overall results regarding the knowledge and practices were arbitrarily graded as very poor, poor, average, good and very good if the percentages of barbers who responded positively ranged from 0 to 20%, 21–40%, 41–60%, 61–80% and 81–100% respectively. Also since we could not observe the practices of barbers directly, we collected the information regarding the practices as reported by them.

2.1. Statistical analysis

The data collected was tabulated, coded and analyzed using Analysis Toolpak in Microsoft Excel. Chi square test and Fishers' exact test were used for evaluating the statistical significance of the association between the independent and the dependent factors. For all the tests, p value <0.05 was considered significant.

3. Results

The mean age group of the barbers was 32.05 ± 11.02 years with majority in the age group 20–40 years and the mean number of years in the profession of barbering was 11.58 ± 9.56 years. Average number of clients at each shop was 35.9 ± 18.4 . Three-fifth of the barbers had an education of high school or more. Ninety percent of the barbers had heard the term HIV; however the correct nature of disease was known to only 12% of the RSB. The corresponding proportions were better in SB. Knowledge regarding all the modes of transmission was miserable with none of the RSB knowing all the modes of transmission. Also, blood contact acting as a mode of transmission was known to only 5.6% of the RSB. Myths regarding the mode of transmission were rampant and contaminated water, mosquito bite and fomites were commonly cited. Knowledge regarding any of the symptoms of HIV was absent in about 45% of the barbers.

Most of the RSB were not aware about the method of diagnosis of HIV. They preferred to visit private and government hospitals for getting themselves diagnosed. However, overall awareness about VCTC (Voluntary Counseling and Testing Center) was poor (20.2%). The overall knowledge regarding the prevention methods related to their profession such as disinfection of the instruments and use of new blades was average. About half of the RSB did not know about any of the prevention methods. Anti-retroviral therapy as a treatment of HIV was known to very few (7.8%).

Most of the SB were aware of the risk of transmission of HIV through their profession, while only one fourth of the RSB knew about it. The knowledge regarding hepatitis was even poorer. Most effective means of information was television followed by person to person communication and radio (Table 1).

All the barbers had movable razors, while one fifth of them also used the fixed razor. While all the SB reportedly provided and used a new blade voluntarily for every customer, the corresponding figures were lower for the RSB. The use of a new blade voluntarily for the side lock and axilla dipped down further in both the categories. Reported disinfection of instruments was average on the whole; however the average contact time was less than a minute. Various unsafe methods of disposal were in practice such as indiscriminate throwing which was most customary in RSB, disposal in public bins which was widespread in SB. Blades were also sold away and even taken away by people and sweepers.

The most common treatment for cuts was application of alum in both the groups. The SB used other materials as well, such as Dettol, Savlon and after shave lotion (Table 2).

4. Discussion

The present study was an attempt to study the knowledge regarding HIV/AIDS among the barbers which are one of the potential high-risk groups due to their exposure to all the sections of the population as well as the increased risk of injuries owing to their profession. In the absence of appropriate disinfection procedures and knowledge regarding prevention methods, the instruments at the barbers shop can serve as a potential mode of transmission of the disease especially during the rush hours when the client crowding occurs.¹⁴

Majority of the barbers were in the economically productive age group which therefore is a matter of concern lest they should contract the disease owing to ignorance. We observed a wide gap in knowledge regarding various aspects of HIV/AIDS in the RSB which could be due to their lower educational status as compared to the SB. However since none of the barbers were illiterate, educational status could not be used as a plea for ignorance, especially with the ongoing campaigns in mass media. This becomes even more relevant in the light of the fact that the average years in the profession of barbering were around 11 years, signifying that they have lived with the existing practices for long. It was quite dispiriting to encounter myths regarding the mode of transmission in both the groups indicating a need for enhanced IEC activities. The deficiency in knowledge regarding the infectious nature of the disease and especially the infectious risks associated with exposure to blood which is important for their occupation was depressing, yet comparable to the results of other studies.^{15,16} However, another study from Ethiopia among the barbers reported better knowledge (78%) regarding the spread of HIV through non-sterile equipments.¹⁷

The unawareness regarding the symptoms of HIV disease makes these high-risk group individuals vulnerable to late reporting and consequently late diagnosis. To make the situation even worse, the knowledge regarding the availability of

Table 1 – Comparison of knowledge regarding HIV in shop barbers and road side barbers in Kanpur.

Knowledge about	Shop barbers (270)		Road side barbers (90)		Total (360)		χ^2 Value, <i>p</i> value
	No.	%	No.	%	No.	%	
Nature of the disease^a							
Infectious	187	69.3	11	12.2	198	55.0	88.72, 0.00
Hereditary	0	0	8	8.9	8	2.2	24.55, 0.00
Other	83	30.7	35	38.9	118	32.8	2.03, 0.15
Don't know	0	0	36	46.7	36	10.0	120.0, 0.00
Mode of transmission^a							
Sexual transmission	235	87.0	19	21.1	254	70.6	141.22, 0.00
Blood contacts	189	70.0	5	5.6	194	53.8	112.82, 0.00
From mother to the child	99	36.7	8	8.9	107	29.7	24.93, 0.00
Blood transfusion	189	70.0	9	10.0	198	55.0	98.18, 0.00
All of the above	18	6.7	0	0	18	5.0	4.99, 0.00
Myths in mode of transmission^a							
Contact	23	6.7	11	12.2	34	9.4	1.08, 0.29
Contaminated water	108	40.0	37	41.1	145	40.2	4.73, 0.02
Air	83	26.7	17	18.9	100	27.8	12.14, 0.00
Mosquito bite	162	60.0	35	38.9	197	54.7	10.44, 0.00
Fomites	21	6.7	18	20.0	39	10.8	17.73, 0.00
Symptoms^a							
Mild illness (sore throat, fatigue, malaise, rash)	111	41.1	15	16.7	126	35.0	17.73, 0.00
Persistent generalized lymphadenopathy	33	12.2	8	8.9	41	11.4	0.74, 0.38
Diarrhea	92	34.1	7	7.8	99	27.5	23.41, 0.00
Loss of weight >10%	2	0.7	0	0	2	0.6	0.00, 1.00
Fever, night sweats, thrush	130	48.1	6	6.7	136	37.8	49.41, 0.00
Others	34	12.6	11	12.2	45	12.5	0.01, 0.92
Don't know	108	40.0	53	58.9	161	44.7	9.74, 0.00
Method of diagnosis							
Lab test	136	50.4	2	2.2	138	38.3	66.2, 0.00
Don't know	134	49.6	88	97.8	222	61.7	
Prevention methods^a							
Condom use	198	73.3	10	11.1	208	57.8	107.13, 0.00
Use of new syringes	180	66.7	4	4.4	184	51.1	104.58, 0.00
Disinfection of the instruments used for shaving	133	49.2	21	23.3	154	42.8	18.53, 0.00
Use of new blade for shaving	165	61.1	53	58.9	218	60.6	0.14, 0.70
Safe blood transfusion	132	40.0	3	3.3	135	37.5	59.77, 0.00
Others	39	14.4	0	0	39	10.8	14.58, 0.00
Don't know	35	13.0	43	47.8	78	21.7	48.21, 0.00
Treatment modalities							
ARV	28	10.4	0	0	28	7.8	10.12, 0.00
Don't know	242	89.6	90	100	332	92.2	
Diseases transmitted through barbering^a							
HBV/HCV	66	24.4	7	7.8	73	20.3	11.6, 0.00
AIDS/HIV	237	87.8	22	24.4	259	71.9	134.14, 0.00
Skin infection	150	55.5	37	41.1	187	51.9	5.64, 0.01
Others	36	13.3	20	22.2	56	15.6	4.06, 0.04
Don't know	37	13.7	0	0	37	10.3	13.75, 0.00
Source of information^a							
TV	226	83.7	25	27.8	251	69.7	115.26, 0.00
Radio	131	48.5	37	41.1	168	46.7	1.49, 0.22
Person to person	200	74.1	9	10.0	209	58.1	113.80, 0.00
Newspaper	163	60.4	18	20.0	181	50.3	44.01, 0.00
Others	38	14.1	5	5.6	43	11.9	4.66, 0.03
a Multiple responses.							

a Multiple responses.

facilities for HIV diagnosis, counseling and treatment at VCTC as well as that of anti-retroviral treatment for the disease was largely deficient. Information regarding the availability of these centers should be distributed to all sections of the society so that they can avail the benefits of the programme.

HIV virus is susceptible to heat and sunlight and survives outside the body for seconds usually but may survive up to 10–15 min^{18,19}

HIV is completely inactivated by treatment for 10 min at room temperature with any of the following: 10% household

Table 2 – Comparison of practices regarding HIV in shop barbers and road side barbers in Kanpur.

Practices	Shop barbers (270)		Road side barbers (90)		Total (360)		χ^2 Value, p value
	No.	%	No.	%	No.	%	
Type of razor^a							
Fixed	51	18.9	36	40.0	87	24.17	16.42, 0.00
Movable	270	100	90	100	360	100.00	–
Supply of new blade voluntarily for every customer	270	100	68	75.6	338	93.89	70.30, 0.00
Use of new blade voluntarily for^a							
Shaving	270	100	76	84.4	346	96.11	39.64, 0.00
Side lock	236	87.4	67	74.4	303	84.17	8.51, 0.00
Axilla	236	87.4	53	58.9	289	80.28	34.67, 0.00
Disinfection of instruments done^a	162	60.0	20	22.2	182	50.56	38.54, 0.00
Beta propiolactone	33	12.2	1	1.1	34	9.44	9.74, 0.001
Ether	29	10.7	0	0.0	29	8.06	10.51, 0.00
Heating	10	3.7	2	2.2	12	3.33	0.11, 0.73
Acetone	21	7.8	0	0	21	5.83	7.43, 0.006
Ethanol	42	15.5	7	7.8	49	13.61	3.47, 0.06
Other (Savlon, Dettol, Soap water, Water etc.)	27	10.0	10	11.1	37	10.28	0.09, 0.76
Contact time (in minutes)	0.67 ± 1.01		0.67 ± 0.48				
Disposal of blades^a							
Indiscriminate	41	15.2	57	63.3	98	27.22	78.98, 0.00
In public bins	146	54.1	6	6.7	152	42.22	62.19, 0.00
Taken by sweeper	58	21.5	8	8.9	66	18.33	7.15, 0.01
Sold away	20	7.4	23	25.6	43	11.94	21.14, 0.00
Taken away by people	18	6.7	7	7.8	25	6.94	0.13, 0.71
Others	19	7.0	0	0	19	5.28	5.35, 0.00
No. of cuts in a week	1 ± 0		2 ± 2				
Material applied on cuts^a							
Alum	255	94.4	71	78.9	326	90.56	19.10, 0.00
Dettol/Savlon	161	59.6	6	6.7	167	46.39	76.13, 0.00
After shave lotion	118	43.7	0	0	118	32.78	58.52, 0.00
Water	54	20.0	13	14.4	67	18.61	1.03, 0.30
Powder	12	4.4	0	0	12	3.33	2.87, 0.04

a Multiple responses.

bleach, 50% ethanol, 35% isopropanol, 1% Nonidet P40, 0.5% Lysol, 0.5% paraformaldehyde, or 0.3% hydrogen peroxide. The virus is also inactivated by extremes of pH (pH 1.0, pH 13.0).²⁰ Half of all the barbers were not disinfecting their instruments and those who did, used beta propiolactone, ether, acetone, ethanol, Savlon, Dettol, soap water and even water for disinfection. However the concentration of the disinfectants used was not known to them and the contact time averaged 0.67 ± 1.01 min and 0.67 ± 0.48 min only, much less than the recommended time. Similar findings in improper disinfection have been reported by other researchers also.^{3,4,16,17,21}

Deficient knowledge regarding the effectiveness of disinfection of barbering instruments as a method for preventing the transmission of HIV increases the risk of transmission of HIV at these shops, both to the barbers as well as the customers. This becomes more important especially during the rush hours when there is a queue of customers, allowing almost no time for effective disinfection.

The most effective way to limit the risk of HIV infection in health care professionals is to use universal precautions with every patient, for example washing hands and wearing protective barriers (gloves, aprons, goggles) and the same stands true for the barbers as well, since the HIV status of the customer

is unknown and the risk of injury always there. This knowledge gap was reflected in their practices as well, with only half of all the barbers practicing disinfection of the instruments. This is especially important in light of the fact that the number of cuts every week averaged from one to two which could be an underreporting since it was not based on an observation rather inquired from the barbers verbally. Workers at other places have reported higher incidences of cuts.^{2–4}

Use of fixed blade and reuse of the same blade are harmful practices adding to the disease risk and have been condemned elsewhere also.^{2,22}

Application of alum on the cuts is another unhygienic practice adding to the risk since often the same piece of alum is rubbed on all the customers. Although the knowledge regarding the transmission risk through a blade was low, the practice was found to be quite high. This discrepancy could be due to the fact that since the clients ask for the use of a new blade owing to the increased awareness programmes, the barbers answered affirmatively on being inquired about the use of a new blade. Yet, the percentages for use of new blade for axilla shaving, which is more prone to injury due to loose skin, were low. Another matter for concern was the various inappropriate methods of disposal of blades, which pose as a risk for transmission of infection to the scavengers and the

general public. This issue has been raised by other researchers as well.²³

The knowledge regarding the risk of transmission of other diseases through the profession of barbering was even lower than that regarding HIV/AIDS. The knowledge regarding HBV and HCV which have a greater potential of spread through the cuts was very poor. Since television and radio were the main source of information, they can be used to increase the awareness. However due emphasis needs to be given to the other methods as well, especially through interpersonal communication which will be fruitful to demystify the myths and queries regarding the disease in barbers. Since barbers are in a unique position of interacting continuously with almost all the male population, they have been used as peer educators in some of the countries and very encouraging results have been obtained.^{24–26}

Replication of the model would provide us good results but would require intensive health education campaigns for education of the barbers.

Although the sexual mode of transmission remains the most common route but the chances of transmission through barbering and various other similar indigenous professions such as tattooing, ceremonial balding of children, delivery by dais etc which involve the risk of exposure to blood should not be overlooked. In-service training that involves the barbers could help us not only prevent HIV transmission but also simultaneously provide a check for the transmission of other more susceptible parenterally transmitted infections such as hepatitis infections. A baseline prevalence of the disease according to the occupation of an individual would guide us to more hidden and underestimated modes of transmission.

4.1. Limitations

The selection of barbers was done non-randomly and therefore generalization of findings is limited although the findings of the study are still important in Indian perspective where studies in similar area are scarce. Also we could not observe the practices of the barbers which could provide us a more realistic picture, due to the paucity of time. Follow up of the study participants which is a limitation of a cross-sectional study was another issue which could provide us a more conclusive evidence of the association of barbering and HIV.

4.2. Strengths and future directions

Very few studies are available from India regarding the knowledge, awareness and practices of barbers regarding HIV, therefore this study addresses a lesser explored avenue of prevention of HIV/AIDS. Contact with contaminated razors leading to HIV and other blood borne infections is a serious concern and needs to be dealt with caution. Correlating the serological status of the barbers with their practices, which was not possible due to ethical and logistic concerns would provide us more evidence for the occupational hazard of the profession. Health education, training, supervision and

monitoring would help in facilitating the adoption of safe practices among the barbers.

Conflicts of interest

All authors have none to declare.

REFERENCES

1. Melkote SR, Goswami D. Predictors of attitudes to persons with AIDS among young adults in India. *Indian J Soc Work*. 2000;61(1):89–105.
2. Khandait DW, Ambadekar NN, Vasudeo ND. Knowledge and practices about HIV transmission among barbers of Nagpur city. *Indian J Med Sci*. 1999;53:167–171.
3. Zewudie T, Legesse W, Kurkura G. Knowledge, attitudes and practices among barbers in South-western Ethiopia. *Afr Newsl Occup Health Saf.*. 2000;1269–1271.
4. Ibrahim MT, Opara WE, Tanimowo T. Knowledge of HIV/AIDS, infection prevention practices and accidental skin cuts in barbing saloons in Sokoto, Nigeria. *Nigeria Med Pract*. 2007;51(6):123–127.
5. Salami KK, Titiloye MA, Brieger WR, Otusanya SA. Observations of barbers' activities in Oyo state, Nigeria: implications for HIV/AIDS transmission. *Int Q Community Health Educ*. 2006;24(4):319–330.
6. Wazir MS, Mehmood S, Ahmed A, Jadoon HR. Awareness among barbers about health hazards associated with their profession. *J Ayub Med Coll Abbottabad*. 2008;20(2):35–38.
7. *Current and Future Dimensions of the HIV/AIDS Pandemic – A Capsule Summary*. Geneva: G.P.A. WHO Publication; 2002:1–15.
8. *Guidelines for the Nursing Management of People with AIDS*. WHO AIDS Series. Geneva: WHO Publication; 1988, 10–21.
9. Donald A. *Modes of Transmission, the Basis for the Prevention Strategies, AIDS Prevention and Control*. Geneva: WHO Publication; 1988:14–19.
10. Marcus R, Kay K, Mann JM. Transmission of HIV in health care setting world wide. *Bull World Health Organ*. 1989;67(5):577.
11. *Guideline on Sterilization and High-level Disinfection Methods Effective Against HIV; AIDS Action Who Report: Global Program on AIDS*. 1988.
12. Gilada IS. AIDS and barbers. *CARC Calling*. 1992;5:33–34.
13. Olivetti M. AIDS and barber shop. *CARC Calling*. 1992;5:32.
14. Kalasagar M, Sivapathasundaram B, Einatein T, Bertin A. AIDS awareness in an Indian metropolitan slum dweller: a KAP (Knowledge, Attitude, Practice) study. *Ind J Dent Res*. 2006;17(2):66–69.
15. Zahraoui-Mehadji M, Baakrim MZ, Laraqui S, et al. Infectious risks associated with blood exposure for traditional barbers and their customers in Morocco. *Sante*. 2004 Oct–Dec;14(4):211–216.
16. Beyen TK, Tulu KT, Abdo AA, Tulu AS. Barbers' knowledge and practice about occupational biological hazards was low in Gondar town, North West Ethiopia. *BMC Public Health*. 2012;12:942.
17. Biadagegn F, Belyhun Y, Anagaw B, et al. Potential risk of HIV transmission in barbering practice in Ethiopia: from public health and microbiological perspectives. *BMC Public Health*. 2012;12:707.
18. Von Reyn CF, Mann JM, Chin J. International travel and HIV infection. *WHO Bull*. 1990;68:251–259.

19. WHO AIDS Series 2. *Guidelines on Sterilization and Disinfection Method Effective Against HIV*. 2nd ed. Geneva: WHO Publication; 1989:2–10.
20. Jawetz, Melnick, Adelberg's AIDS & Lentiviruses. In: Brooks GF, Carroll KC, Butel JS, Morse SA, eds. *Medical Microbiology*. McGraw Hill's Company; 2007.
21. Arulogun Oyedunni S, Adesoro Moses O. Potential risk of HIV transmission in barbering practice among professional barbers in Ibadan, Nigeria. *Afr Health Sci*. Mar 2009;9(1):19–25.
22. Khaliq AA, Smego RA. Barber shaving and blood-borne disease transmission in developing countries. *S Afr Med J*. 2005;95(2):94–96.
23. Janjua NZ, Nizamy MAM. Knowledge and practices of barbers about hepatitis B and C transmission in Rawalpindi and Islamabad. *J Pak Med Assoc*. Mar 2004;54(3):116–119.
24. *The Nation's Health*. Black Community Groups Confront HIV/AIDS Threat: Eliminating Racial, Ethnic Disparities.
25. John JK. Country watch: India. *AIDS STD Health Promot Exch*. 1995;3:7–8.
26. Howard J. Raising a chorus of voices to prevent HIV. *Harv AIDS Rev*. 19952–3.

Copyright © 2012, INDIACLEN. Publishing Services by Reed Elsevier India Pvt Ltd. All rights reserved.

<http://dx.doi.org/10.1016/j.cegh.2012.11.003>

Original Article

Alcohol use, HIV risk behavior and experience of sexually transmitted infections among female sex workers of Nepal

Reshu Agrawal Sagtani^{a,*}, Sailesh Bhattarai^b, Baikuntha Raj Adhikari^c, Dharanidhar Baral^d, Deepak Kumar Yadav^e, Paras Kumar Pokharel^f

^a Senior Resident, School of Public Health and Community Medicine, B.P. Koirala Institute of Health Sciences, Dharan, Nepal

^b Assistant Professor, School of Public Health and Community Medicine, B.P. Koirala Institute of Health Sciences, Dharan, Nepal

^c Assistant Professor, Department of Psychiatry, B.P. Koirala Institute of Health Sciences, Dharan, Nepal

^d Assistant Professor, School of Public Health and Community Medicine, B.P. Koirala Institute of Health Sciences, Dharan, Nepal

^e Associate Professor, School of Public Health and Community Medicine, B.P. Koirala Institute of Health Sciences, Dharan, Nepal

^f Professor, Chief, School of Public Health and Community Medicine, B.P. Koirala Institute of Health Sciences, Dharan, Nepal

ARTICLE INFO

Article history:

Received 18 March 2013

Accepted 22 May 2013

Available online 6 June 2013

Keywords:

Alcohol

Condom

Risk behavior

ABSTRACT

Background: Alcohol use is a pervasive phenomenon among female sex workers and their clients. It affects their physical health and promotes high-risk behavior and consequently, sexually transmitted infections (STIs). Hence, we designed a study to find the factors associated with alcohol use among the female sex workers of Eastern Nepal. The prevalence of HIV high-risk behavior and STIs among sex workers was also recorded.

Methods: A cross sectional study was conducted in three commercial districts namely Sunsari, Morang and Jhapa of Eastern Nepal. All the FSWs who had been involved in commercial sex activity in the past six months were included in the study. Face-to-face interviews with 168-restaurant based and 42-street based female sex workers, who were identified through snowball sampling. Bivariate analyses were conducted using Chi-Square tests for categorical variables. Multivariate models were used to identify significant factors associated with alcohol use in the sex trade.

Results: Logistic regression analyses revealed that sex with intravenous drug user (IDUs) (AOR 2.275, 95% CI 1.266, 16.898), oral sex (AOR 3.261, 95% CI 1.166–9.121) and pregnancy

* Corresponding author. Tel.: +977 9842051828.

E-mail address: rsagtani@hotmail.com (R.A. Sagtani).