Available online at www.sciencedirect.com

ScienceDirect

journal homepage: www.elsevier.com/locate/cegh

Original Article

Blood component therapy in neonates in a neonatal intensive care unit of northern India

Avneet Kaur^a, Shashi Kant Dhir^{b,*}, Gurmeet Kaur^c, Manish Gupta^a,
Meenal Batta^d

^a Post Graduate Student, Department of Pediatrics, Guru Gobind Singh Medical College, Faridkot, Punjab, India

^b Assistant Professor, Department of Pediatrics, Guru Gobind Singh Medical College, Faridkot, Punjab, India

^c Professor and Head, Department of Pediatrics, Guru Gobind Singh Medical College, Faridkot, Punjab, India

^d Senior Resident, Department of Physiology, Guru Gobind Singh Medical College, Faridkot, Punjab, India

ARTICLE INFO

Article history:

Received 15 September 2015

Accepted 30 October 2015

Available online 26 November 2015

Keywords:

Blood component therapy

Transfusion

Neonates

ABSTRACT

Background: Neonates, especially preterm babies admitted in neonatal intensive care units (NICU), frequently require transfusion of blood components. The use of blood components in the neonates varies across different hospitals. There is urgent need of forming global guidelines and policies for blood component transfusions in the sick neonates. Although the guidelines for Indian setup from the premier institutes exist, yet there is paucity of Indian data regarding the actual blood component usage.

Aims and objective: To know the incidence and indications of various blood components transfusion in our NICU.

Material and method: This retrospective single center observational study was done in a tertiary care NICU of a government medical college of northern India. The records of all neonates admitted in the NICU from August 2014 to July 2015 were screened for blood component usage and their indications. The data were statistically analyzed using Microsoft Excel™ software.

Results: During the study period, 815 neonates were admitted in NICU of our hospital. Out of these neonates, 280 (34.3%) neonates received at least single blood component transfusion at some point of NICU stay. More than one type of components was needed in 100 (12.2%) neonates. A total of 557 units of blood components were transfused in the babies. The mean blood component transfusion was 1.98 ± 0.21 in the babies requiring transfusion. The maximum (54.7%) component to be transfused was platelets followed by packed red blood cells (24.5%). Sepsis was the most common reason for blood component transfusion.

Conclusion: A large number of admitted neonates require blood components. Sepsis remains the foremost etiological factor for blood component transfusion.

© 2015 INDIACLEN. Published by Elsevier, a division of Reed Elsevier India, Pvt. Ltd. All rights reserved.

* Corresponding author. Tel.: +91 9780190130.

E-mail address: drshashikantdhir@gmail.com (S.K. Dhir).

<http://dx.doi.org/10.1016/j.cegh.2015.10.011>

2213-3984/© 2015 INDIACLEN. Published by Elsevier, a division of Reed Elsevier India, Pvt. Ltd. All rights reserved.

1. Introduction

Neonates admitted in neonatal intensive care units (NICUs) require transfusion of blood and blood components for various reasons. With the advent of newer techniques, the sick neonates stay for more time in NICUs and they are one of the most common recipients of whole blood (WB) or blood components. Gone are the days when only WB was used. From last quarter century, blood components are used almost exclusively to decrease avoidable transfusion reactions, sensitization, volume overload, and to use one unit of blood to save many lives. Various blood components include packed red blood cells (PRBC), fresh frozen plasma (FFP), and platelet concentrate (PC). Preterm babies comprise major group requiring transfusion, and 85% of extremely low birth weight babies admitted in NICUs have required blood component transfusion at some point of time.¹

The common reasons in the neonatal intensive care unit for which various blood components are transfused include thrombocytopenia, anemia, bleeding, various surgical conditions, exchange transfusion for hyperbilirubinemia, etc. Thrombocytopenia is common in sick neonates admitted in NICU. Platelet transfusions are life saving for bleeding manifestations accompanied with thrombocytopenia. Most platelet transfusions are still given prophylactically to thrombocytopenic neonates even without bleeding.² Previous studies suggest that neonatal thrombocytopenia is a risk factor for hemorrhage (particularly intra-ventricular hemorrhage), mortality, and adverse neurodevelopmental outcome.³ Thrombocytopenic neonates who receive platelets are up to ten times more likely to die than neonates who do not receive platelet transfusion. Platelet transfusions are given to all neonates with platelet counts below 30,000/dl and below 100,000/dl in other circumstances.⁴ FFP is transfused in neonates with significant coagulopathy. Common coagulopathies in neonates include vitamin K deficiency and disseminated intravascular coagulation due to sepsis. Packed RBC transfusion is also common transfusion given in NICU to treat anemia depending upon the comorbid conditions as per the guidelines. Exchange transfusions using WB are done in hyperbilirubinemia and sepsis.

The use of blood components in the neonates varies across different hospitals. There is urgent need of forming global guidelines and policies for blood component transfusions in the sick neonates. Although the guidelines for Indian setup from the premier institutes exist, yet there is paucity of Indian data regarding the actual blood component usage.⁵ This study was conducted to know the incidence of transfusion of various blood components in our NICU and various indications for which they are being used. Although western data are available on usage of isolated blood components, there is paucity of Indian data for the blood component usage. This study was planned to get an insight into above-mentioned issues.

2. Material and methods

This retrospective single center observational study was done in a tertiary care neonatal intensive care unit of a government

medical college of northern India. The records of all neonates admitted in the NICU from August 2014 to July 2015 were screened for the blood component usage and the indications of the transfusion. All consecutive patients younger than 28 days admitted during the study period in NICU were eligible for study enrollment. The neonatal intensive care unit of our hospital caters to the neonates suffering from various critical diseases from adjacent districts and states. All the inborn, outborn, and babies admitted with surgical causes were included. The neonates, who stayed in NICU for less than twenty-four hours, received any blood component prior to admission, or who were readmitted, were excluded from the study. Our hospital has certified blood bank which provides WB and its various fractions, i.e. PRBC, FFP, PC, and platelet-rich plasma. Facility for cryoprecipitate is also available at our blood bank. The indications of transfusing various blood components being followed in our NICU are detailed in [Table 1](#). Random donor platelets were used for the transfusion.

The details of PRBC, FFP, PC, and WB transfusions were recorded from the files. The mean value of hemoglobin, platelet count, and coagulogram indices was noted at the time of concerned blood component transfusion. The mean age of the neonates at the time of transfusion was noted. Detailed information about the components transfused, number of transfusions, and indications was noted in a predesigned validated Proforma. The data were taken till the final outcome of the baby. The data were statistically analyzed using Microsoft Excel™ software.

3. Results

During the study period, 815 neonates were admitted in NICU of our hospital. The baseline variables of the neonates included in the study are shown in [Table 2](#). Out of enrolled neonates, 280 (34.3%) neonates received at least single blood component transfusion at some point of NICU stay. More than one type of components was needed in 100 (12.2%) neonates. The number of units of blood components transfused in the babies was 557. The mean blood component transfusion was 1.98 ± 0.21 in the babies requiring transfusion. The component-wise detail of various transfusions is being shown in [Table 3](#). The various etiologies of need of blood component transfusion are listed in [Table 4](#).

The incidence of thrombocytopenia (platelet count less than 1.5 Lakhs/cmm) was 42.9% in neonates. The incidence of severe (less than 0.5 Lakhs), mild (1–1.5 Lakhs/cmm), and moderate thrombocytopenia (0.5–1 Lakhs/cmm) was 49.5%, 28.5% and 22% respectively. Mean hemoglobin at the time of transfusion of packed red blood cell was 8.2 ± 1.3 g/dl. The mean platelet count for transfusing the platelets was $35,458 \pm 3,962$ /dl. The mean prothrombin time index (PTI) was $63.8 \pm 6\%$ and partial thromboplastin time (PTT) was 55.2 ± 6.2 s in the babies requiring transfusion of FFP. The mean age of platelet transfusion was 5.2 ± 1.1 days and the overall mean age of blood component transfusion in the neonates in days was 13 ± 1.63 .

In the multiple transfusions group, the random donor platelets were given more than once in 51% of the transfused

Table 1 – Transfusion guidelines used in our NICU.

Component	Indication
Packed red blood cell	<ul style="list-style-type: none"> Asymptomatic infants with Hct <21% and reticulocytes <100,000/μL (2%) Infants with Hct <31% and hood O₂ <36% or mean airway pressure <6 cm H₂O by CPAP or IMV or >9 apneic and bradycardic episodes per 12 h or 2/24 h requiring bag and mask ventilation while on adequate methylxanthine therapy or HR >180/min or RR >80/min sustained for 24 h or weight gain of <10 g/day for 4 days on 100 kcal/kg/day or having surgery Infants with Hct <36% and requiring >35% O₂ or mean airway pressure 6–8 cm H₂O by CPAP or IMV
Platelet	<p><30,000/cubic mm</p> <ul style="list-style-type: none"> Transfuse all <p>30–49,000/cubic mm</p> <ul style="list-style-type: none"> BW <1500 g and $\leq$7 days old Clinically unstable Concurrent coagulopathy Previous significant hemorrhage (grade 3 or 4 IVH) Prior to surgical procedure Post-operative period (72 h) <p>50,000–100,000/cubic mm</p> <ul style="list-style-type: none"> Active bleeding NAIT with intracranial bleed Before or after neurosurgical procedures
Fresh frozen plasma	<ul style="list-style-type: none"> Disseminated intravascular coagulation Deranged coagulogram Prothrombin time index >2.0 Partial thromboplastin time >55s in preterm Partial thromboplastin time >50s in term
Whole blood	<ul style="list-style-type: none"> Double volume exchange transfusion

Hct, Hematocrit; CPAP, Continuous positive airway pressure; IMV, Intermittent Mechanical Ventilation; BW, birth weight; IVH, Intra-ventricular hemorrhage; NAIT Neonatal alloimmune thrombocytopenia.

neonates. Regarding second transfusion of similar type, the platelets were transfused maximum number of times (33.5%) followed by FFP (12.5%). None of the babies admitted during study period required cryoprecipitate. No transfusion

reaction was seen in any neonate during the whole study period.

4. Discussion

This study was done to see the pattern of blood component usage and reasons for the transfusion in a NICU of northern India to get an insight of the existing transfusion practices.

Our study showed more use of blood components as compared to the WB which is similar to the studies done elsewhere.^{1,6} The most common component administered was platelets followed by red cells and this pattern has also been found in earlier studies.⁶ For the FFP transfusion, results similar to our study both in indications and frequency were seen in a multicenter study done in Italy.⁷ Although the current evidence does not support treating the abnormal indices, this practice is being followed.^{3,7} The reason of giving platelets was sepsis akin to the studies done by the both Asian and western people.^{8,9} Sepsis per se remains most common cause or associated factor in blood component therapy and almost all of the babies requiring multiple types of blood components had sepsis as primary diagnosis.

Despite a lot of research on the blood component therapies, a consensus is still to be made on various threshold values for transfusions of blood components. Answers to queries like optimal platelet transfusion threshold optimal red blood cell (RBC) transfusion threshold and new practices to prevent

Table 2 – Baseline variables.

Variable	Value
Males	432 (53%)
Birth weight in grams	1265 $\pm$ 134
Gestation by last menstrual period	33 weeks and 1 day
Gestation by New Ballard Score ¹³	32 weeks and 5 days
Preterm	538 (66%)
Mode of delivery	
Unassisted vaginal delivery	423 (52%)
Cesarean-section with labor	41 (5%)
Cesarean section without labor	310 (38%)
Breech vaginal	41 (5%)
Perinatal score	3 (3–3)
Duration of rupture of membranes (h)	14.21 $\pm$ 45.12
Duration of labor (h)	5.69 $\pm$ 1.46
Apgar at 1 min	8 (7–9)
Apgar at 5 min	9 (8–9)
Length of hospital stay in days	1.02 $\pm$ 1.23

Values represent mean standard deviation, median (interquartile range). Percentages are given in parenthesis.
New Ballard Score¹³ was used for gestation assessment in all babies as per our unit protocols.

Table 3 – Details of individual blood component transfusion.

Component	Number of neonates	Total units of transfusions	Once	Twice	Thrice	>Thrice
PRBC	115 (30.2)	137 (24.5)	101 (38.2)	9 (11.5)	5 (25)	0 (0)
WB	12 (3.1)	15 (2.6)	10 (3.8)	1 (1.2)	1 (5)	0 (0)
PC	173 (45.5)	305 (54.7)	88 (33.3)	58 (74.3)	9 (45)	18 (100)
FFP	80 (21.0)	100 (17.9)	65 (24.6)	10 (12.8)	5 (25)	0 (0)

Values in parenthesis indicate percentages.

PRBC, packed red blood cells; WB, whole blood; PC, platelet concentrate; FFP, fresh frozen plasma.

Table 4 – Etiology of blood component transfusion.

Blood component	Indication	Number/percentage
Platelet concentrate	Sepsis	125/72.2
	HIE	20/11.5
	MAS	9/5.2
	PM	13/7.5
	Surgical	6/3.4
Fresh frozen plasma	Active bleeding	50/62.5
	Deranged coagulogram	30/37.5
Packed red blood cell	Sepsis	57/49.5
	HIE	8/6.9
	Surgery	24/20.8
	PM	20/17.3
	Others	6/5.2
Whole blood	Neonatal jaundice	9/75
	Sepsis	3/25

HIE, hypoxic ischemic encephalopathy; MAS, meconium aspiration syndrome; PM, prematurity.

transfusion-transmitted infections are still not clear but further research would help to answer them.¹⁰ Evidence-based guidelines are still sidelined to the expert opinion and efforts are required to bring them to practice.¹¹ A large-scale randomized multicenter controlled trial (PlaNet-2 trial) will probably be able to give more precise guidelines about the thresholds required to transfuse platelets.¹²

The limitations of our study were small sample size, retrospective nature, and single center in our study. Another limitation of the study was not keeping track of the volume of blood drawn for various samplings during the stay. A prospective stratified study with different gestations and birth weights with a fairly large sample size will help us in acquiring more knowledge about the blood component transfusion practices.

5. Conclusion

Blood components rather than WB are frequently used in NICUs. Sepsis remains the foremost etiological factor for blood component transfusion, especially if there is need for multiple transfusions. Abnormal coagulograms are still being treated with FFPs. Although a lot of data exist regarding the transfusion practices in the neonates, further studies are still required to fill the gaps.

What this study adds

- Sepsis is the most common reason for blood component transfusion, especially in patients requiring multiple transfusions.
- Platelet transfusions are most frequently required in neonates.
- Abnormal coagulograms are still being treated.

Conflicts of interest

The authors have none to declare.

Acknowledgements

We are thankful to Dr Jasmeen, Dr Veerinder, Dr Aastha, and Dr Saminder for their help in the data collection and filling the proformas. We are also thankful to medical record department of our hospital for providing us the patient's files. We are thankful to Dr R.K. Goyal for his continuous support and proof reading the final manuscript.

REFERENCES

1. Jain R, Jose B, Coshic P, Agarwal R, Deorari AK. Blood and blood component therapy in neonates. *Indian J Pediatr.* 2008;75(May (5)):489–495.
2. Christensen RD, Carroll PD, Josephson CD. Evidence-based advances in transfusion practice in neonatal intensive care units. *Neonatology.* 2014;106(3):245–253.
3. Murray NA, Roberts IAG. Neonatal transfusion practice. *Arch Dis Child Fetal Neonatal Ed.* 2004;89(March (2)):F101–F107.
4. Cloherty JP, Eichenwald EC, Hansen AR, Stark AR, eds. In: *Manual of Neonatal Care* 7th ed. USA: Lippincott William & Wilkins; 2012:585.
5. Agarwal R, Deorari A, Paul VK, eds. In: *AIIMS Protocols in Neonatology* 1st ed. New Delhi: CBS Publishers & Distributors; 2015.
6. Giridharan J, Ramalingam GR. Study of blood component therapy in neonates. *Int J Sci Res.* 2015;4(March (3)):196–200.
7. Motta M, Del Vecchio A, Perrone B, Ghirardello S, Radicioni M. Fresh frozen plasma use in the NICU: a prospective,

- observational, multicentred study. *Arch Dis Child Fetal Neonatal Ed.* 2014;99(July (4)):F303–F308.
8. Del Vecchio A. Evaluation and management of thrombocytopenic neonates in the intensive care unit. *Early Hum Dev.* 2014;90(Suppl. 2):S51–S55.
 9. Gupta AK, Kumari S, Singhal A, Bahl A. Neonatal thrombocytopenia and platelets transfusion. *Asian J Transfus Sci.* 2012;6(July (2)):161–164.
 10. Nickel RS, Josephson CD. Neonatal transfusion medicine: five major unanswered research questions for the twenty-first century. *Clin Perinatol.* 2015;42(September (3)):499–513.
 11. Parker RI. Transfusion in critically ill children: indications, risks, and challenges. *Crit Care Med.* 2014;42(March (3)):675–690.
 12. Curley A, Venkatesh V, Stanworth S, et al. Platelets for neonatal transfusion – Study 2: A randomised controlled trial to compare two different platelet count thresholds for prophylactic platelet transfusion to preterm neonates. *Neonatology.* 2014;106(2):102–106.
 13. Ballard JL, Khoury JC, Wedig K, Wang L, Eilers-Walsman BL, Lipp R. New Ballard score, expanded to include extremely premature infants. *J Pediatr.* 1991;119(September (3)):417–423.