Available online at www.sciencedirect.com

ScienceDirect

journal homepage: www.elsevier.com/locate/cegh

Original Article

An enumeration of the prevalence of hypothyroidism during pregnancy in central India

Arindam Bose^a, Neelu Soni^b, Nivadita Dashore^b, Kamal Gajria^b,
Sushma Jhamad^b, Nanda Hemvani^a, Dhananjay S. Chitnis^{a,*}

^a Department of Pathology, Choithram Hospital and Research Centre, Indore, India

^b Department of Obstetrics and Gynaecology, Choithram Hospital and Research Centre, Indore, India

ARTICLE INFO

Article history:

Received 31 August 2015

Accepted 13 November 2015

Available online 2 December 2015

Keywords:

Hypothyroidism

Pregnancy

First trimester

Central India

Thyroid stimulating hormone (TSH)

ABSTRACT

Background/objective: Hypothyroidism can have adverse effects on pregnancy, such as miscarriage and congenital hypothyroidism. The routine assessment of thyroid function is not being done to date on all the pregnant women in India. Hence, the present study was designed to survey the antenatal thyroid status during the first trimester of pregnancy in a central Indian population.

Methods: The present study was performed on 1152 pregnant women with a mean age of 26.25 ± 4.39 years. All women were in the first trimester and were screened for their thyroid stimulating hormone (TSH) and Free T₄ level.

Results: Following the guidelines for the TSH and Free T₄ values in pregnancy, the prevalence of the hypothyroidism among the pregnant women in central India was 13.1%. Furthermore, the prevalence of clinical hypothyroid and subclinical hypothyroid was 2.25% and 10.85%, respectively.

Conclusion: The present data on the prevalence of hypothyroidism in pregnant women strongly recommend the inclusion of routine thyroid screening.

© 2015 INDIACLEN. Published by Elsevier, a division of Reed Elsevier India, Pvt. Ltd. All rights reserved.

1. Introduction

Thyroid disorders are the second most common endocrinopathies found during pregnancy. The prevalence of hypothyroidism during pregnancy is estimated to be 0.3–0.5% for overt hypothyroidism and 2–3% for subclinical hypothyroidism. Thyroid autoantibodies are found in 5–15% of women in the childbearing age and are a risk factor for hypothyroidism

during pregnancy and the postpartum period. Hashimoto thyroiditis is the most common cause.¹ A population survey conducted in 1991 showed that 2.5% of pregnant women have compensated hypothyroidism based on elevated thyroid stimulating hormone (TSH) levels.² A meta-analysis of 10 prospective studies of euthyroid women revealed that women with antithyroid antibodies were more than twice likely to experience a miscarriage (odds ratio 2.30, 95% CI 1.80–2.95).³

* Corresponding author. Tel.: +91 731 2362491–99; fax: +91 731 2470068; mobile: +91 9302124512.

E-mail address: ds_chitnis@rediffmail.com (D.S. Chitnis).

<http://dx.doi.org/10.1016/j.cegh.2015.11.004>

2213-3984/© 2015 INDIACLEN. Published by Elsevier, a division of Reed Elsevier India, Pvt. Ltd. All rights reserved.

There are very limited studies from India on thyroid disorders in pregnancy. Some studies based on Indian population showed the insufficiency of iodine in pregnant women.⁴ A study from Mumbai, India, highlighted on the optimization of thyroid management in pregnancy and the improved neonatal outcomes associated with it.⁵

However, in many developing countries, routine assessment of thyroid status is not being done in all pregnant women. Furthermore, the incidence of subclinical hypothyroidism or autoimmune status in this population of women, especially women in South Asia or India is not properly enumerated due to lack of data.

The thyroid plays a key role in the fetal neural development, so the mother's thyroid status has to be within normal range during pregnancy. The present study was aimed to survey antenatal thyroid status during the first trimester of pregnancy in central India.

2. Materials and methods

The present study included 1152 pregnant women, who turned up from different parts of the central India to the Out Patient Department of Choithram Hospital and Research Centre, Indore, India, between February 2009 and March 2013. In this study, 257 (22.31%) subjects were from low income group ($\leq$ Rs.10,000 per month), 343 (29.77%) were from high income group ($\geq$ Rs. 40,000 per month) and the majority of the subjects, 552 (47.92%) were from middle income group ($>$ Rs. 10,000 to $<$ Rs. 40,000 per month). The study subjects were between 18 and 39 years with the mean age of 26.25 ($\pm$ 4.39) years. Known cases of thyroid disorders were excluded from this study. The study proposal was reviewed by the hospital Ethical Committee, which has guidelines based on Helsinki declaration.

All the subjects visited at the Out Patient Department (OPD) of Choithram Hospital and Research Centre, Indore, India, during the first trimester of their pregnancy, were included in this study; and screened for their TSH and Free T₄ levels. Other than biochemical thyroid assessment, pregnancy profile included hemoglobin, urea, creatinine, random blood glucose, HIV, HBsAg, Rubella Antibody, and routine urine examination.

Pelvic examination and ultrasound examination were done. Details of pregnancy regarding gestational age, weight gain and menstrual history were recorded. Features suggestive of hypothyroidism, such as family history of thyroid disorders, previously diagnosed thyroid disease, presence of thyroid therapy, skin changes associated with hypothyroidism (cold skin, coarse skin), presence of goiter, slow movements, periorbital puffiness and delayed ankle reflex were looked for.

General physical examination including height and weight measurements was also done for all. Body mass index (BMI) was calculated for all the subjects. The normal range of BMI lies between 18.5 and 24.9 kg/m². BMI range between 24.9 kg/m² and 30 kg/m² were considered as overweight, whereas more than 30 kg/m² was considered as obese.⁶

TSH and Free T₄ were assayed using Abbott Architect 1100SR (Abbott Laboratories, Diagnostics Division, Abbott

Park, IL, USA) analyzer based on a Chemiluminescent Microparticle Immunoassay (CMIA) for the quantitative determination of TSH (Architect TSH assay was designed to have a precision of $\leq$ 10% total CV and the assay was designed to have a functional sensitivity of $\leq$ 0.01 μ LU/ml, which meets the requirements of a third generation TSH assay) and Free T₄ (The Architect Free T₄ assay was designed to have a precision of $\leq$ 10% total CV and the assay was designed to have an analytical sensitivity of $\leq$ 0.4 ng/dl).⁷ The guidelines for the pregnant women, suggested that TSH should be less than 2.5 μ IU/ml in the first trimester and hence was considered as cut off value for TSH in the present study.⁸ Whereas the normal range of the Free T₄ was considered as 0.7–1.48 ng/dl.

The data collected were analyzed using Excel 2007, R2.8.0 Statistical Package for Social Sciences (SPSS) for windows version 16.0 (SPSS Inc.; Chicago, IL, USA).

3. Results

Out of total 1152 subjects screened, 1101 (95.58%) had serum TSH levels within normal range with a mean TSH value of 1.96 ± 1.2 μ IU/ml. On the contrary 26 subjects (2.25% of the total study population) had clinical hypothyroidism with a higher mean TSH value of 26.87 ± 6.51 μ IU/ml with decreased value of Free T₄ with the mean of 0.53 ± 0.03 ng/dl and 125 subjects (10.85% of the total study population) had a mean TSH value of 7.43 ± 1.2 μ IU/ml and Free T₄ values in normal range (mean value of 0.94 ± 0.16 ng/dl), considered as subclinical hypothyroidism (Fig. 1).

Out of these 26 clinical hypothyroid subjects, 20 (76.9%) were obese and rest of 6 (33.9%) were overweight. On the other

Fig. 1 – A graphical presentation of the thyroid status during first trimester of pregnancy in central India (normal = 1001, clinical hypothyroidism = 26 and subclinical hypothyroidism = 125).

side, out of 125 subclinical hypothyroid subjects, 38 (30.4%) were obese, 76 (60.8%) were overweight, and rest of the 11 (8.8%) subjects had normal BMI.

4. Discussion

The association between thyroid disorders and adverse pregnancy outcomes has been studied earlier in western countries.⁹ Pop et al. reported that maternal hypothyroidism and hypothyroxinemia occurring in the first trimester of pregnancy may be harmful to the embryo-fetal brain development and lead to intellectual retardation in the offspring.¹⁰ Data on the prevalence of hypothyroidism among pregnant women in Indian population appear scanty.^{11,12}

The present study demonstrates a higher incidence of hypothyroidism. The prevalence of overall hypothyroidism in this cohort is higher than that in the western literature; the prevalence ranges from 2.6%¹³ to 4.8%¹⁴ and a previous Indian study (3.69%).¹² However, a study conducted in 2013 on Indian population showed 14.3% women had hypothyroidism and the majority of these women had subclinical hypothyroidism.¹⁵ The recent review on undiagnosed subclinical hypothyroidism also showed the prevalence of the disease in pregnant women in the range of 3–15%.¹⁶

In the present study, 125 (10.85%) subjects were having subclinical hypothyroidism and the prevalence is much higher compared to western data.^{13,14} Among the cases of hypothyroidism (26 clinical hypothyroid and 125 subclinical hypothyroid), only 38 had undergone TPOAb assessment (The ARCHITECT Anti-TPO assay was designed to have an assay precision of <10% total CV for samples >5.61 IU/mL and the assay was designed to have an analytical sensitivity of <1.0 IU/mL) and the results showed that 31 out of 38 had elevated level of serum TPOAb.

The benefits and risks of universal testing or case-finding strategies in pregnant women are debated.¹⁷ Brent had previously pointed out that the potential adverse outcomes are so significant, and the tools to diagnose and treat thyroid disease are easily accessible that it is no longer acceptable for even one third of pregnant women to remain undiagnosed.¹⁸ Abalovich et al. showed that untreated hypothyroidism, subclinical, or overt, at the time of conception is associated with miscarriage rate of 31.4% compared with 4% in euthyroid subjects at conception.⁹

USG thyroid and antithyroid antibody tests were not done for the majority of the hypothyroid cases in the present study, and that could have provided additional information. The present data indicate a high prevalence of hypothyroidism among pregnant women in the population and supports performing universal screening early in pregnancy for thyroid disorders using the first trimester-specific reference intervals. The prevalence of stillbirth, premature delivery, and/or other complications during delivery were not reported in the hypothyroid patient population at our end, probably due to the adequate treatment given to the patients to maintain euthyroid state. Based on the observation of the present study we strongly recommended TSH measurement

as the initial screening test for thyroid dysfunction in the first trimester.

5. Conclusion

This study concludes that the prevalence of overall hypothyroidism during pregnancy in central India is too high in contrast to western countries. The present study strongly recommends the adaptation of universal thyroid screening for all the pregnant women during the first trimester in Indian population.

Conflicts of interest

The authors have none to declare.

Acknowledgements

We express our gratitude to the management of Choithram Hospital and Research Centre, Indore for providing infrastructure and facilities. We gratefully acknowledge all the resident gynecologists and nurses for their valuable tips and kind cooperation.

REFERENCES

1. Abalovich M, Amino N, Barbour LA, et al. Management of thyroid dysfunction during pregnancy and postpartum: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab.* 2007;92(8):S1–S47.
2. Klein RZ, Haddow JE, Faix JD, et al. Prevalence of thyroid deficiency in pregnant women. *Clin Endocrinol (Oxf).* 1991;35:41–46.
3. Prummel MF, Wiersinga WM. Thyroid autoimmunity and miscarriage. *Eur J Endocrinol.* 2004;150:751–755.
4. Pathak P, Singh P, Kapil U, Raghuvanshi RS. Prevalence of iron, vitamin A, and iodine deficiencies amongst adolescent pregnant mothers. *Indian J Pediatr.* 2003;70:299–301.
5. Nuzhat A, Pranathi R, Evita F. Hypothyroidism in pregnancy: is universal screening needed. *J Obstet Gynecol India.* 2006;56:495–498.
6. Committee on Obstetric Practice: The American College of Obstetricians and Gynecologist. *Weight Gain during Pregnancy.* Committee Opinion No. 548. 2013 (Reaffirmed 2015): 1–3.
7. Kalani M, Arachchige K, Hadlow N, Wardrop R, Lim EM, John P. Age-specific TSH reference ranges have minimal impact on the diagnosis of thyroid dysfunction. *Clin Endocrinol.* 2012;77(5):773–779.
8. Mazzaferrri EL. Editors' Choice – TSH levels in pregnancy: TSH levels between 2.5 and 5.0 in first-trimester thyroid antibody-negative women are associated with a significant increase in the rate of spontaneous miscarriage. *Clin Thyroid.* 2010;22(7):6–8.
9. Abalovich M, Gutierrez S, Alcaraz G, Maccallini G, Garcia A, Levalle O. Overt and subclinical hypothyroidism complicating pregnancy. *Thyroid.* 2002;12:63–68. <http://dx.doi.org/10.1089/10507250.2753451986>.

10. Pop VJ, Brouwers EP, Vader HL, Vulmsa T, van Baar AL, de Vijlder JJ. Maternal hypothyroxinaemia during early pregnancy and subsequent child development: a 3-year follow-up study. *Clin Endocrinol*. 2003;59:282–288. <http://dx.doi.org/10.1046/j.1365-2265.2003.01822.x>.
11. Sahu MT, Das V, Mittal S, Agarwal A, Sahu M. Overt and subclinical thyroid dysfunction among Indian pregnant women and its effect on maternal and fetal outcome. *Arch Gynecol Obstet*. 2010;281(2):215–220.
12. Mukhopadhyay A, Pati S, Mukherjee S, Das N, Mukhopadhyay P, Saumandal B. Autoimmune thyroid disorders and pregnancy outcome: a prospective observational study. *Thyroid Res Pract*. 2007;4(1):50–52.
13. Vaidya B, Anthony S, Bilous M, et al. Detection of thyroid dysfunction in early pregnancy: universal screening or targeted high-risk case finding? *J Clin Endocrinol Metab*. 2007;92(1):203–207.
14. Nambiar V, Jagtap VS, Sarathi V, et al. Prevalence and impact of thyroid disorders on maternal outcome in Asian-Indian pregnant women. *J Thyroid Res*. 2011;1–6. <http://dx.doi.org/10.4061/2011/429097>.
15. Dhanwal DK, Prasad S, Agarwal AK, Dixit V, Banerjee AK. High prevalence of subclinical hypothyroidism during first trimester of pregnancy in North India. *Indian J Endocrinol Metab*. 2013;17(2):281–284.
16. Negro R, Stagnaro-Green A. Diagnosis and management of subclinical hypothyroidism in pregnancy. *BMJ*. 2014;349:g4929.
17. Surks MI, Ortiz E, Daniels GH, et al. Subclinical thyroid disease: scientific review and guidelines for diagnosis and management. *JAMA*. 2004;291:228–238. <http://dx.doi.org/10.1001/jama.291.2.228>.
18. Brent GA. Diagnosing thyroid dysfunction in pregnant women: is case finding enough? *J Clin Endocrinol Metab*. 2007;92:39–41. <http://dx.doi.org/10.1210/jc.2006-2461>.