Letter to the editor

Platelets and community acquired pneumonia

We have greatly enjoyed reading the recently published article by Jain et al.¹ In this prospective observational study, the authors included 152 children between 2–59 months admitted with severe pneumonia. This study revealed that cyanosis, severe acute malnutrition, platelet count < 0.7 lac/mm³ on admission predicted mortality in patients with community-acquired pneumonia (CAP).

Community-acquired pneumonia is one of the leading causes of hospitalization, in-hospital death and long-term mortality. The prevalence of thrombocytopenia at hospital admission ranged from 2% to 5% among consecutive adult CAP patients to 25% in CAP patients admitted at the intensive care units.² On the other hand, thrombocytosis on admission has been observed among 8–13% of CAP inpatients.² The presence of both thrombocytopenia and thrombocytosis at hospital admission was associated with poor short-term outcome in adult patients with CAP.² However, platelet count may vary significantly within several days or weeks. A decrease in platelet count over time is usually caused by reduced production, increased destruction, and/or sequestration of thrombocytes and a rapid rise in platelet count reflects increased platelet formation by megakaryocytes, indicating adequate compensatory response of bone marrow to various stimuli including infection and inflammation.³ Therefore, we think that Jain and colleagues should be evaluated not only the prognostic significance of low platelet counts but also the significance of increased platelet counts, mean platelet volume, and time-dependent platelet count changes in children with CAP.

References

1. Jain A, Awasthi N, Awasthi S. Low platelet counts predict mortality in severe community acquired pneumonia in children under 5 years of age: a hospital based observational study. *Clin Epidemiol Glob Health*. 2018. <http://dx.doi.org/10.1016/j.cegh.2018.02.007>.
2. Mirsaeidi M, Peyrani P, Aliberti S, et al. Thrombocytopenia and thrombocytosis at time of hospitalization predict mortality in patients with community-acquired pneumonia. *Chest*. 2010;137(February (2)):416–420.
3. Gorelik O, Izhakian S, Barchel D, et al. Prognostic significance of platelet count changes during hospitalization for community-acquired pneumonia. *Platelets*. 2017;28(June (4)):380–386. <http://dx.doi.org/10.1080/09537104.2016.1219032> Epub 2016 Sep 29.

Oğuzhan Çelik*, Cem Çil

Muğla University, Faculty of Medicine, Department of Cardiology, Turkey
E-mail address: oguzhan.celik1@saglik.gov.tr

Funda Sungur Biteker

Yatağan State Hospital, Department of Infectious Diseases and Clinical Microbiology, Turkey

Eda Özlek

Muğla University, Faculty of Medicine, Department of Cardiology, Turkey

Bülent Özlek

Muğla University, Faculty of Medicine, Department of Cardiology, Turkey

Volkan Doğan, Murat Biteker

Muğla University, Faculty of Medicine, Department of Cardiology, Turkey

* Corresponding author at: Mugla Sıtkı Kocman Üniversitesi Tıp Fakültesi, Kotekli Mahallesi, Marmaris Yolu, No:48, 48000, Muğla, Turkey.

<https://doi.org/10.1016/j.cegh.2018.06.001>

Received 6 May 2018

Available online 05 June 2018

2213-3984/ © 2018 Published by Elsevier, a division of RELX India, Pvt. Ltd on behalf of INDIACLEN.