Validation of the Hamilton Anxiety Rating Scale and State Trait Anxiety Inventory A and B in Arabic among the Lebanese population

Souheil Hallit^{a,b,*}, Chadia Haddad^{a,1}, Rabi Hallit^a, Marwan Akel^{b,d}, Sahar Obeid^{a,e,f}, Georges Haddad^{a,c}, Michel Soufia^a, Wael Khansa^a, Rony Khoury^a, Nelly Kheir^g, Christiane Abi Elias Hallit^c, Pascale Salameh^{b,h,i}

^a Faculty of Medicine and Medical Sciences, Holy Spirit University of Kaslik (USEK), Jounieh, Lebanon

^b INSPECT-LB: Institut National de Santé Publique, Epidémiologie Clinique et Toxicologie, Beirut, Lebanon

^c Psychiatric Hospital of the Cross, Jal Eddib, Lebanon

^d School of Pharmacy, Lebanese International University, Beirut, Lebanon

^e Faculty of Philosophy and Human Sciences, Holy Spirit University, Kaslik, Lebanon

^f Faculty of Pedagogy, Lebanese University, Beirut, Lebanon

^g Faculty of Pedagogy, Université de la Sainte Famille, Batroun, Lebanon

^h Faculty of Pharmacy, Lebanese University, Beirut, Lebanon

ⁱ Faculty of Medicine, Lebanese University, Beirut, Lebanon

ARTICLE INFO

Keywords:

Anxiety
HAM-A
STAI-A
STAI-B
Arabic
Validation

ABSTRACT

Objectives: Our aim in this study was to translate the Hamilton Anxiety Rating Scale (HAM-A), State Trait Anxiety Inventory (STAI-A and B) scales into Arabic, linguistically validate them for use in a representative sample of the Lebanese population, and to check the reliability of these Arabic versions.

Methods: The study was cross-sectional, conducted between November 2017 and March 2018, which enrolled 1332 community-dwelling adults, using a proportionate random sample from all Lebanese Mohafazat.

Results: The factor analyses of the anxiety scales were run over the whole sample (N = 1332). All of the HAM-A, STAI-A and STAI-B items could be extracted from the list. All items from all the scales did not over-correlate to each other (r < 0.9), did not have a low loading on factors (< 0.3) or a low communality (< 0.3). The factor analyses showed 2 factors for HAM-A (Cronbach alpha = 0.921), 3 factors for STAI-A (Cronbach alpha = 0.928) and 2 factors for STAI-B (Cronbach alpha = 0.898). A significantly high ICC was found between the HAM-A, STAI-A (ICC = 0.766) and STAI-B (0.704). In addition, a significantly high ICC was found between the STAI-A and STAI-B scales (ICC = 0.884).

Conclusion: The linguistically validated Arabic versions of these scales can be used to screen for anxiety among the Lebanese populations.

1. Introduction

Anxiety is a transient emotional state affected by individual personality traits that varies across culture and situations.¹ The intensity of anxiety varies over time and differs from person to person according to stressful life events, situations and cultural background.² It is considered as a universal phenomenon that can be present across cultures.

The severity of anxiety symptoms could be measured by using different psychometric tools such as the state-trait anxiety inventory (STAI) and the Hamilton Anxiety Rating Scale (HAM-A). These scales

are widely and extensively used in research and clinical studies and are available in different versions for adults and children.^{3–6} The STAI has been validated and translated to several languages and is considered as a standard international tool for measuring anxiety in research.⁵ A validated Arabic version of the STAI, with an adequate internal consistency reliability, exists among patients attending dental clinics.⁷ Also, the HAM-A has been proven to have a high reliability and sensitivity for measuring anxiety and has been validated among adults and adolescents as well.^{8,9}

More specifically, the **Hamilton Anxiety Rating Scale (HAM-A)** is a psychological questionnaire used by clinicians to rate the severity of a

* Corresponding author. Building 560, Street 8, Biakout, Lebanon.

E-mail address: souheilhallit@hotmail.com (S. Hallit).

¹ First co-authors.

patient's anxiety. Anxiety can denote things such as “a mental state ... a drive ... a response to a particular situation ... a personality trait ... and a psychiatric disorder”.¹⁰ Nonetheless it was one of the first anxiety rating scales to be published, the HAM-A remains widely used by clinicians.⁹ It was originally published by Max Hamilton in 1959. For clinical purposes, only severe or inadequate anxiety is attended to. This scale is considered a “clinical rating” of the range of anxiety, and is proposed for individuals that are “already diagnosed with anxiety neurosis”.¹¹

As for the **State-Trait Anxiety Inventory (STAI)**, it measures two types of anxiety – state anxiety, or anxiety about an event, and trait anxiety, or anxiety level as a personal characteristic; and used in diagnoses, in both clinical and other medical settings.⁶ The advantage of this scale is that could be applied towards assessing different types of anxiety. This was a new development because all other questionnaires focused on one type of anxiety at the time.¹²

Appropriate, validated and translated scales are needed to explain and predict anxiety in each country due to the influence of linguistic and cultural differences. The use of the validated scale in different languages, to assess the severity of the anxiety symptoms, demands two concepts: (1) maintaining a level of psychometric properties comparable to the original version and (2) adapting the scale to the cultural of the country. The validated and translated scale with a good reliability, will be helpful in clinical settings and research.¹³

Therefore, it is essential to validate the Arabic version of the STAI and HAM-A scales for cultural adaptation among the Lebanese population in order to measure the severity of anxiety. Our aim in this study was to translate the HAM-A, STAI-A and B scales to Arabic, linguistically validate them for use in a representative sample of the Lebanese population, and check the reliability of these Arabic versions.

2. Methods

2.1. Study design and sampling (Sample 1)

This study is cross-sectional, conducted between November 2017 and March 2018, which enrolled 1332 community-dwelling participants using a proportionate random sampling from the 15 Lebanese Mohafazat (Beirut, Mount Lebanon, North South and Bekaa). Each Mohafaza is divided into Caza (stratum) two villages were randomly selected from the list of villages provided by the Central Agency of Statistics in Lebanon. Patients were randomly selected from each village. All participants above 18 years old were eligible to participate. Excluded were the patients with psychiatric problems, mental retardation, dementia or who refused to answer the questionnaire. Data collection was performed through personal interviews with participants by trained, study independent personnel.

2.2. Ethical approval

The Psychiatric Hospital of the Cross Ethics and Research Committee, in compliance with the Hospital's Regulatory Research Protocol, approved this study protocol (HPC-009-2018) based on the fact that the autonomy and confidentiality of participants were respected and since it was an observational study, no harm will be prompted to them. The purpose and requirement of the study was informed to each patients. Consent was obtained as written approval on the ethical consent form.

2.3. Minimal sample size calculation

Comrey and Lee suggested that a minimum of 10 observations per variable is necessary in order to avoid computational difficulties.¹⁴ Since the STAI-A and B scales questionnaire contains 40 questions (20 questions for each scale), a minimal sample of 400 patients was needed to conduct an exploratory factor analysis.

Table 1
Sociodemographic characteristics of the sample population.

	Frequency (%)
Gender	
Male	480 (36.1%)
Female	848 (63.9%)
Education level	
Illiterate	13 (1.0%)
Primary	21 (1.6%)
Complementary	64 (4.8%)
Secondary	224 (16.9%)
University	883 (66.7%)
Higher education	119 (9.0%)
Employment status	
Unemployed	917 (68.8%)
Employed	415 (31.2%)
Socioeconomic status	
< 1000\$	926 (73.5%)
1000–2000 \$	251 (19.9%)
> 2000 \$	83 (6.6%)
Marital status	
Single	963 (72.6%)
Married	334 (25.2%)
Widowed	13 (1.0%)
Divorced	17 (1.3%)
Father's education level	
Illiterate	68 (5.1%)
Primary	162 (12.2%)
Complementary	279 (21.0%)
Secondary	335 (25.3%)
University	482 (36.3%)
Mother's education level	
Illiterate	96 (7.3%)
Primary	127 (9.6%)
Complementary	302 (22.9%)
Secondary	393 (29.8%)
University	402 (30.5%)
Residence	
Alone	95 (7.2%)
With others	1217 (92.8%)
Personal history of medical illness	
Yes	157 (11.8%)
No	1174 (88.2%)
Family history of psychiatric illness	
Yes	69 (5.2%)
No	1262 (94.8%)
Mean ± SD	
Age (in years)	28.08 ± 16.79
Number of kids	1.15 ± 1.72

2.4. Questionnaire

The questionnaire used during the interviewed was in Arabic, the native language of Lebanon. A trained staff was in charge of collecting the data, via a personal interview with each participant. This person was independent of this study. The first part assessed the socio-demographic characteristics of the included population (age, gender, educational level of the patient and his parents, marital status, socioeconomic level, alcohol consumption, family history of psychiatric diseases). The socioeconomic level, defined as the family monthly income, was divided into 3 categories: low (< 1000 USD, intermediate (1000–2000 USD) and high (2000 USD). Also, we asked the patient if he got a physician diagnosis of anxiety and was taking a medication to treat it. The other parts comprised the different scales used in this study as follows:

2.5. Hamilton anxiety scale (HAM-A)

The HAM-A¹⁵ entails 14 items, each categorized by a series of symptoms, and measures mental agitation and psychological distress, as

Table 2

Promax rotated matrix of HAMA scale factors, Sample 1, and confirmatory factor analysis parameters, Sample 2 (italics).

	Items	Factor 1	Factor 2
Genitourinary symptoms	12	0.881/0.401	
Cardiovascular symptoms	9	0.821/0.584	
Autonomic symptoms	13	0.800/0.742	
Respiratory symptoms	10	0.774/0.633	
Gastrointestinal symptoms	11	0.721/0.597	
Somatic (sensory)	8	0.667/0.810	
Behavior at interview	14	0.643/0.376	
Somatic (muscular)	7	0.551/0.694	
Tension	2		0.910/0.916
Anxious mood	1		0.903/0.842
Depressed mood	6		0.748/0.942
Insomnia	4		0.692/0.756
Fears	3		0.529/0.521
Intellectual	5		0.486/0.596

Factor 1 = Somatic items; Factor 2 = Psychic/Psychological items.

Cronbach's alphas: factor 1 = 0.898; factor 2 = 0.853; total scale = 0.921.

Percentage of variance explained: 58.51%.

Kaiser-Meyer-Olkin (KMO) = 0.947.

Bartlett's test of sphericity = $p < 0.001$.

well as anxiety-related physical complaints. The responses on the scale were measured on a 5-point Likert scale: 0 (symptoms not present), 1 (mild symptoms), 2 (moderate symptoms), 3 (severe symptoms) and 4 (very severe symptoms). The total score was calculated by summation of the 14 items.

2.6. State-Trait Anxiety Inventory (STAI) A and B

It is composed of 40 questions that measure two types of anxiety: state anxiety, or anxiety about an event (STAI-A), and trait anxiety, or anxiety level as a personal characteristic (STAI-B). The answers follow a 4-point Likert scale, with 1 (not at all), 2 (somewhat), 3 (moderately so) and 4 (very much so) for STAI-A and 1 (almost never), 2 (sometimes), 3 (often) and 4 (almost always) for STAI-B. Higher scores are positively correlated with higher levels of anxiety.¹⁶

Table 3

Promax rotated matrix of STAI-A scale factors, Sample 1, and confirmatory factor analysis parameters, Sample 2 (italics).

	Items	Factor 1	Factor 2	Factor 3
I feel steady	19	0.795/0.497		
I feel pleasant	20	0.713/0.635		
I feel content	16	0.696/0.609		
I feel self-confident	11	0.659/0.557		
I feel satisfied	8	0.597/0.668		
I feel at ease	5	0.582/0.612		
I feel comfortable	10	0.538/0.580		
I am relaxed	15	0.467/0.570		
I am tense	3		0.779/0.694	
I feel strained	4		0.705/0.544	
I feel nervous	12		0.647/0.833	
I am worried	17		0.601/0.747	
I feel upset	6		0.587/0.765	
I feel calm	1		0.510/0.609	
I feel secure	2		0.508/0.488	
I feel frightened	9			0.721/0.564
I am jittery	13			0.706/0.583
I feel indecisive	14			0.635/0.671
I am presently worrying over possible misfortunes	7			0.586/0.533
I feel confused	18			0.449/0.283

Cronbach's alphas: factor 1 = 0.852; factor 2 = 0.855; factor 3 = 0.780; total scale = 0.928.

Percentage of variance explained: 56.04%.

Kaiser-Meyer-Olkin (KMO) = 0.953.

Bartlett's test of sphericity = $p < 0.001$.

2.7. Forward translation into Arabic

A single bilingual translator, Arabic native fluent in English, aware of the concepts of the three anxiety scales translated the English versions of the scales into Arabic. An expert committee, composed of health care professionals (psychiatrists and psychologists), a language professional and the original translator, had reviewed and revised the translated questionnaire in order to check for idiomatic and conceptual equivalence of the Arabic translated version.^{17–20}

2.8. Back translation into English

A native English speaker translator person, fluent in Arabic, had back translated the Arabic versions of the three scales into the English language. The translator was unaware with the concepts of the anxiety scales and the original English versions.^{17,18} The expert committee compared the back-translated English version of the questionnaire with the original English one, in order to check for inconsistencies and to solve any discrepancies between the versions. All ambiguities disappeared after repeating the process of forward-back translation.^{17–22}

2.9. Sample 2

We conducted a further cross-sectional study in May 2018 on a sample of Lebanese patients attending psychology clinic that enrolled 155 participants. To ensure the validity of the results, the scales were tested on another sample (sample 2), independent from the first one. Patients filled a questionnaire through a face-to-face interview.

Statistical analyses

The SPSS software version 23 was used for data analysis. A descriptive analysis were done using the counts and percentages for categorical variables and mean and standard deviation for continuous measures. The anxiety scales validity was tested in two different methods. First, construct validity was determined by an exploratory factor analysis, using the principal components analysis technique (Sample 1): The Bartlett's test of sphericity and Kaiser-Meyer-Olkin measure of sampling adequacy were confirmed to be adequate. The number of factors retained corresponded to Eigenvalues higher than

Table 4

Promax rotated matrix of STAI-B scale factors, Sample 1, and confirmatory factor analysis parameters, Sample 2 (italics).

	Items	Factor 1	Factor 2	Factor 3	Factor 4
I am happy	30	0.827/0.693			
I feel pleasant	21	0.818/0.607			
I feel rested	26	0.815/0.675			
I am content	36	0.779/0.749			
I feel satisfied with myself	23	0.726/0.736			
I feel secure	33	0.721/0.693			
I am a steady person	39	0.656/0.561			
I make decision easily	34	0.511/0.547			
I am calm, cool, and collected	27	0.489/0.328			
Some unimportant thoughts runs my mind and bothers me	37		0.807/0.642		
I get in a state of tension or turmoil as I think over my recent concerns and interests	40		0.729/0.618		
I take disappointments so keenly that I can't put them out of my mind	38		0.709/0.585		
I worry too much over something that really doesn't matter	29		0.707/0.701		
I have disturbing thoughts	31		0.678/0.628		
I feel nervous and restless	22		0.626/0.528		
I feel that difficulties are piling up so that I cannot overcome them	28		0.455/0.328		
I feel inadequate	35			0.755/0.585	
I lack self-confidence	32			0.700/0.597	
I feel like a failure	25			0.680/0.681	
I wish I could be as happy as others seem to be	24				0.898/0.413

Cronbach's alphas: factor 1 = 0.875; factor 2 = 0.839; factor 3 = 0.742; total scale = 0.898.

Percentage of variance explained: 58.27%.

Kaiser-Meyer-Olkin (KMO) = 0.940.

Bartlett's test of sphericity = $p < 0.001$.**Table 5**

Intraclass correlation coefficient between insomnia scales.

		HAMA	STAI-A	STAI-B
HAMA	ICC (CI)	N/A	0.709 (0.676–0.739)	0.704 (0.671–0.734)
	p-value		< 0.001	< 0.001
STAI-A	ICC (CI)	0.709 (0.676–0.739)	N/A	0.884 (0.871–0.896)
	p-value	< 0.001		< 0.001
STAI-B	ICC (CI)	0.704 (0.671–0.734)	0.884 (0.871–0.896)	N/A
	p-value	< 0.001	< 0.001	

ICC: Intraclass Correlation coefficient; CI: Confidence interval.

one. Items with factor loading > 0.4 are considered as belonging to a factor. The Cronbach's alpha values and the Intraclass correlation coefficient were used to evaluate the internal consistency of the scales. An ICC > 0.7 indicate a good reproducibility.²³

Second, a confirmatory factor analysis was carried out in Sample 2 using the maximum likelihood method for discrepancy function to assess the structure of the structure. We reported several goodness-of-fit indicators: the Relative Chi square (χ^2/df), the Root Mean Square Error of Approximation (RMSEA), the Goodness of Fit Index (GFI) and the Adjusted Goodness of Fit Index (AGFI). The value of χ^2 divided by the degrees of freedom (χ^2/df) has a low sensitivity to sample size and may be used as an index of goodness of fit (cut-off values: $< 2-5$). The RMSEA tests the fit of the model to the covariance matrix. As a guideline, values of < 0.05 indicate a close fit and values below 0.11 an acceptable fit. The GFI and AGFI are chi-square-based calculations independent of degrees of freedom. The recommended thresholds for acceptable values are ≥ 0.90 .²⁴

3. Results

The sociodemographic characteristics of the participants are summarized in Table 1. The results showed that the mean age of the participants was 28.08 ± 16.79 years, with 63.9% females. The majority (75.7%) had a university level of education, unemployed (68.8%), single (72.6%), with a low monthly income (< 1000 USD) (73.5%).

Fig. 1. ROC curve of the HAM-A scale. Patients diagnosed with anxiety by the physician were analyzed. Area under the curve = 0.795 [0.760–0.830] ($P < 0.001$); at value = 11.50, Se = 76.3% and Sp = 71.1%.

Only 5.2% of the participants had a family history of psychiatric illnesses, whereas 11.8% had a history of medical illness.

3.1. Factor analysis

Three factor analyses for the anxiety scales were run over the whole sample ($N = 1332$). All of the HAM-A, STAI-A and STAI-B items could be extracted from the list. All items from all the scales did not over-correlate to each other ($r > 0.9$), did not have a low loading on factors (< 0.3) or a low communality (a communality is the extent to which an item correlates with all other items) (< 0.3). The number of loading factors, KMO and Bartlett's test of sphericity values and the Cronbach alpha values for each scale are summarized in Table 2 (HAM-A), Table 3

Fig. 2. ROC curve of the STAI-A scale. Patients diagnosed with anxiety by the physician were analyzed. Area under the curve = 0.771 [0.732–0.810] ($P < 0.001$); at value = 42.50, Se = 71.3% and Sp = 69.1%.

Fig. 3. ROC curve of the STAI-B scale. Patients diagnosed with anxiety by the physician were analyzed. Area under the curve = 0.779 [0.741–0.817] ($P < 0.001$); at value = 45.50, Se = 73.1% and Sp = 71.6%.

(STAI-A) and Table 4 (STAI-B) respectively.

3.2. Sample 2

A confirmatory factor analysis was run on sample 2, using the structure obtained in Sample 1. The following results were obtained: for the HAMA scale, the Maximum Likelihood Chi-Square = 152.509 and Degrees of Freedom = 76, which gave an $\chi^2/df = 2.01$. For non-centrality fit indices, the Steiger-Lind RMSEA was 0.077 [0.057–0.096]. Moreover, the Joreskog GFI equaled 0.98 and AGFI equaled 0.93.

For the STAI-A scale the Maximum Likelihood Chi-

Square = 405.995 and Degrees of Freedom = 167, which gave an $\chi^2/df = 2.43$. For non-centrality fit indices, the Steiger-Lind RMSEA was 0.094 [0.082–0.108]. Moreover, the Joreskog GFI equaled 0.789 and AGFI equaled 0.734.

For the STAI-B scale the Maximum Likelihood Chi-Square = 344.374 and Degrees of Freedom = 165, which gave an $\chi^2/df = 2.08$. For non-centrality fit indices, the Steiger-Lind RMSEA was 0.087 [0.074–0.099]. Moreover, the Joreskog GFI equaled 0.806 and AGFI equaled 0.753.

3.3. Intraclass correlation coefficient between insomnia scales

A significantly high ICC was found between the HAM-A, STAI-A (ICC = 0.709) and STAI-B (0.704). In addition, a significantly high ICC was found between the STAI-A and B scales (ICC = 0.884) (Table 5).

3.4. ROC curves

The receiver operating characteristic (ROC) curve of the anxiety score as calculated by the HAM-A scale, comparing patients with a physician diagnosis of anxiety to healthy ones, showed that the area under the curve was high = 0.795 [0.60–0.830] ($P < 0.001$); at value = 11.50, the sensitivity was 76.3% and the specificity was 71.1% (Fig. 1).

The ROC curve of the anxiety score as calculated by the STAI-A scale, comparing patients with a physician diagnosis of anxiety to healthy ones, showed that the area under the curve was high = 0.771 [0.732–0.810] ($P < 0.001$); at value = 42.50, the sensitivity was 71.3% and the specificity was 69.1% (Fig. 2).

The ROC curve of the anxiety score as calculated by the STAI-B scale, comparing patients with a physician diagnosis of anxiety to healthy ones, showed that the area under the curve was high = 0.779 [0.741–0.817] ($P < 0.001$); at value = 42.50, the sensitivity was 73.1% and the specificity was 71.6% (Fig. 3).

3.5. Face validity

When comparing the scales scores between patients with and without physician diagnosis of anxiety, the results showed that significantly higher means HAM-A, STAI-A and B scores were found in patients with a physician diagnosis of anxiety compared to those without the diagnosis ($p < 0.001$ for the three scales) (Fig. 4).

4. Discussion

The Arabic versions of the HAM-A, STAI-A and B scales were validated in the current study for use among the Lebanese population. The results showed a high reliability and validity of these scales that provide an initial evidence that could be used as a screening instrument for anxiety in Lebanon. The translation process was conducted by independent translators, the point that decreases the subjectivity of the scales' adaptation. A comparison of the fit of the form obtained through the confirmatory factor analysis revealed that the 2 factors for HAM-A, 3 factors for STAI-A and 4 factors for STAI-B had high fit indices.

4.1. Validation of the scales

The construct validity of the STAI-A and B scales were adequate because items converged over three factors for STAI-A and 4 factors for STAI-B, with adequate factor loadings for all items. The internal consistency of both scales was similar to that obtained in other translated versions for state (Cronbach alpha of 0.93 for state and 0.92 for trait scales in Greece²⁵ and 0.93 to 0.95 for state and 0.91 to 0.93 for trait anxiety in a group of Lebanese and American students²⁶), but lower to that obtained in. The results of the Arabic version of the STAI validity and reliability, had been similar to the psychometric properties

Fig. 4. Comparison of scales scores between patients with and without physician diagnosis of anxiety.

reported by other studies.^{25,27,28}

Factor analysis indicated that a three-factor for STAI-A and a four-factor for STAI-B provide the best explaining criteria of anxiety enclosing the state and traits subscales. Different models were detected in other studies; Abdullatif²⁶ had found two and three factors for STAI-A and STAI-B for American and Lebanese samples. In a Korean study,²⁷ factor analysis had revealed two factors, while a single factor and 2 factor models was revealed by the confirmatory factor analysis. These variations might be due to cultural differences between countries. In fact, the experience and expression of emotions differ from person to other due to the influence of cultural background.¹³ The physiology of the illness syndromes and social context depend on the cultural beliefs that varied across person, thus anxiety disorders are strongly related to racial, ethnic and cultural factors.²⁹

The construct validity of the HAM-A scale was also adequate because items converged over two factors (Anxiety and Psychological), with adequate factor loadings for all items, consistent to the results obtained from previous research for adults.³⁰ The internal consistency of the HAM-A scale was excellent (0.921), suggesting that this scale's items are able to screen for anxiety among the Lebanese population in an excellent way.

4.2. Validity

The construct validity of the Arabic version of these scales was addressed by calculating sensitivity and specificity of the scale. This study is the first to our knowledge to assess this property during the validation process of these scales. We obtained good areas under the curves, sensitivities and specificities for all three scales. These good results may be due to the fact that we compared used the physician diagnosis variable to compare between individuals with anxiety and those without.

4.3. Limitations

The current study has several limitations. An information bias could have occurred since the patients might not have understood the questions well or over/underestimated the answers to some questions. A selection bias is possible because of the refusal rate. The levels of experiential and expressive anxiety are gender different thus our results could be affected by the unequal number of males and females. In addition, the majority of the participants were young according to the mean age, therefore, future studies should be targeted towards older adults. Despite these limitations, the sample size is acceptable and the

study results can be extrapolated to the whole population because of the study design used.

5. Conclusion

Language barriers are an important obstacle to proper medical communication and management, translation of the HAM-A, STAI-A and B scales into Arabic and validation of the Arabic forms was essential. Linguistically validated Arabic versions of these scales can be used to screen for anxiety among the Lebanese populations. Further studies are needed in other Arabic-speaking countries to confirm our findings and to extrapolate the use of the Arabic versions of these scales.

Conflicts of interest

The authors have nothing to disclose.

Funding sources

None.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.cegh.2019.02.002>.

References

1. Spielberger CD. Anxiety as an emotional state. In: Spielberger CD, ed. *Anxiety: Current Trends in Theory and Research*. vol. 1. New York: Academic Press; 1972:23–49. <https://doi.org/10.1016/B978-0-12-657401-2.50009-5>.
2. Ollendick Thomas, King Neville, Yule William. *International Handbook of Phobic and Anxiety Disorders in Children and Adolescents*. New York: Plenum Press; 1994.
3. Julian LJ. Measures of anxiety: state-trait anxiety inventory (STAI), Beck anxiety inventory (BAI), and Hospital anxiety and Depression scale-anxiety (HADS-A). *Arthritis Care Res*. 2011;63(S11):S467–S472.
4. Spielberger CD. Assessment of state and trait anxiety: conceptual and methodological issues. *South Psychol*. 1985;2(4):6–16.
5. Spielberger CD. *State-Trait Anxiety Inventory: Bibliography*. 2nd ed. Palo Alto: Consulting Psychologists Press; 1989.
6. Spielberger CD, Gorsuch RL, Lushene R, Vagg PR, Jacobs GA. *Manual for the State-Trait Anxiety Inventory*. Palo Alto, CA: Consulting Psychologists Press; 1983.
7. Bahammam MA. Validity and reliability of an Arabic version of the state-trait anxiety inventory in a Saudi dental setting. *Saudi Med J*. 2016;37(6):668.
8. Clark DB, Donovan JE. Reliability and validity of the Hamilton anxiety rating scale in an adolescent sample. *J Am Acad Child Adolesc Psychiatry*. 1994;33(3):354–360.
9. Maier W, Buller R, Philipp M, Heuser I. The Hamilton Anxiety Scale: reliability, validity and sensitivity to change in anxiety and depressive disorders. *J Affect Disord*. 1988;14(1):61–68.

10. McDowell I. *Measuring Health: A Guide to Rating Scales and Questionnaires*. USA: Oxford University Press; 2006.
11. Hamilton M. The assessment of anxiety states by rating. *Br J Med Psychol*. 1959;32(1):50–55.
12. Tilton S. Review of the state-trait anxiety inventory (STAI). *New Notes*. 2008;48(2):1–3.
13. Hofmann SG, Hinton DE. Cross-cultural aspects of anxiety disorders. *Curr Psychiatr Rep*. 2014;16(6):450.
14. Comrey AL, Lee HB. *A First Course in Factor analysis*. Psychology Press; 2013.
15. Hamilton M. The assessment of anxiety states by rating. *Br J Med Psychol*. 1959;32(1):50–55.
16. Spielberger CD, Sydeman SJ. State-Trait Anxiety Inventory and State-Trait Anger expression inventory. In: Maruish ME, ed. *The Use of Psychological Testing for Treatment Planning and Outcome Assessment*. Hillsdale, NJ, USA: L. Erlbaum Associates; 1994:292–321 <https://doi.org/10.1002/9780470479216.corpsy0943>.
17. Beaton DE, Bombardier C, Guillemin F, Ferraz MB. Guidelines for the process of cross-cultural adaptation of self-report measures. *Spine (Phila Pa 1976)*. 2000;25(24):3186–3191.
18. Beaton D, Bombardier C, Guillemin F, Ferraz MB. *Recommendations for the Cross-Cultural Adaptation of Health Status Measures*. New York: American Academy of Orthopaedic Surgeons; 2002:1–9.
19. Al-Bannay H, Jarus T, Jongbloed L, Yazigi M, Dean E. Culture as a variable in health research: perspectives and caveats. *Health Promot Int*. 2014;29(3):549–557.
20. Gutiérrez-Sánchez D, Leiva-Santos JP, Sánchez-Hernández R, Hernández-Marrero D, Cuesta-Vargas AI. Spanish modified version of the palliative care outcome scale-symptoms renal: cross-cultural adaptation and validation. *BMC Nephrol*. 2016 Nov 18;17(1):180.
21. Maneesriwongul W, Dixon JK. Instrument translation process: a methods review. *J Adv Nurs*. 2004;48(2):175–186.
22. Nejari C, El Fakir S, Bendahhou K, et al. Translation and validation of European organization for research and treatment of cancer quality of life Questionnaire -C30 into Moroccan version for cancer patients in Morocco. *BMC Res Notes*. 2014;7:228.
23. Terwee CB, Bot SD, de Boer MR, et al. Quality criteria were proposed for measurement properties of health status questionnaires. *J Clin Epidemiol*. 2007;60(1):34–42.
24. Marsh Herb, Hau KT, Wen Z. In search of golden rules: comment on hypothesis-testing approaches to setting cutoff values for fit indexes and dangers in over-generalizing Hu and Bentler's (1999) findings. *Struct Equ Model A Multidiscip j*. January 2004;11:320–341.
25. Fountoulakis KN, Papadopoulou M, Kleanthous S, et al. Reliability and psychometric properties of the Greek translation of the State-Trait Anxiety Inventory form Y: preliminary data. *Ann Gen Psychiatr*. 2006;5:2.
26. Abdullatif QA. *Adaptation of the State Trait anxiety Inventory in Arabic: A Comparison with the American STAI USA*: University of South Florida; 2004 <http://www.scholarcommons.usf.edu/cgi/viewcontent.cgi?article=1932&context=etd>. Available from:.
27. Bieling PJ, Antony MM, Swinson RP. The State-Trait Anxiety Inventory, Trait version: structure and content re-examined. *Behav Res Ther*. 1998 Jul-Aug;36(7-8):777–788.
28. Barnes LL, Harp D, Jung WS. Reliability and utilization of scores on the Spielberger state-trait anxiety inventory. *Educ Psychol Meas*. 2002;62(4):603–618. <https://doi.org/10.1177/0013164402062005>.
29. Hinton DE. Multicultural challenges in the delivery of anxiety treatment. *Depress Anxiety*. 2012;29(1):1–3.
30. Hofmann SG, Anu Aspinwall MA, Hinton DE. Cultural aspects in social anxiety and social anxiety disorder. *Depress Anxiety*. 2010;25(10):1117–1127.

REMOVED